

# Impairment of intestinal glutathione synthesis in patients with inflammatory bowel disease

B Sido, V Hack, A Hochlehnert, H Lipps, C Herfarth, W Dröge

## Abstract

**Background**—Reactive oxygen species contribute to tissue injury in inflammatory bowel disease (IBD). The tripeptide glutathione (GSH) is the most important intracellular antioxidant.

**Aims**—To investigate constituent amino acid plasma levels and the GSH redox status in different compartments in IBD with emphasis on intestinal GSH synthesis in Crohn's disease.

**Methods**—Precursor amino acid levels were analysed in plasma and intestinal mucosa. Reduced (rGSH) and oxidised glutathione (GSSG) were determined enzymatically in peripheral blood mononuclear cells (PBMC), red blood cells (RBC), muscle, and in non-inflamed and inflamed ileum mucosa. Mucosal enzyme activity of  $\gamma$ -glutamylcysteine synthetase ( $\gamma$ GCS) and  $\gamma$ -glutamyl transferase ( $\gamma$ GT) was analysed. Blood of healthy subjects and normal mucosa from a bowel segment resected for tumour growth were used as controls.

**Results**—Abnormally low plasma cysteine and cystine levels were associated with inflammation in IBD ( $p < 10^{-4}$ ). Decreased rGSH levels were demonstrated in non-inflamed mucosa ( $p < 0.01$ ) and inflamed mucosa ( $p = 10^{-6}$ ) in patients with IBD, while GSSG increased with inflammation ( $p = 0.007$ ) compared with controls. Enzyme activity of  $\gamma$ GCS was reduced in non-inflamed mucosa ( $p < 0.01$ ) and, along with  $\gamma$ GT, in inflamed mucosa ( $p < 10^{-4}$ ). The GSH content was unchanged in PBMC, RBC, and muscle.

**Conclusions**—Decreased activity of key enzymes involved in GSH synthesis accompanied by a decreased availability of cyst(e)ine for GSH synthesis contribute to mucosal GSH deficiency in IBD. As the impaired mucosal antioxidative capacity may further promote oxidative damage, GSH deficiency might be a target for therapeutic intervention in IBD.

(Gut 1998;42:485-492)

Keywords: Crohn's disease; ulcerative colitis; glutathione; amino acids;  $\gamma$ -glutamylcysteine synthetase; mucosa

Infiltrating macrophages and neutrophils are abundantly present in inflamed gut in patients with inflammatory bowel disease (IBD). There is a growing body of evidence that these cells expose the inflamed intestine to substantial oxidative stress by production of reactive

oxygen species (ROS).<sup>1-4</sup> Excessive production of ROS could be also demonstrated for circulating phagocytic cells in patients with IBD<sup>5</sup> and was shown to be involved in several models of experimental colitis.<sup>6-8</sup> The mucosa is endowed with various endogenous antioxidant defence systems to remove ROS resulting from normal metabolism. However, enzyme activities of catalase, superoxide dismutase, and glutathione peroxidase were reported to be low at only 4%, 8%, and 45 %, respectively, in healthy colonic mucosa compared with the liver.<sup>9</sup> ROS are highly toxic to cells and oxygen radical formation in excess of physiological amounts may overtax the limited intestinal antioxidant defence system initiating oxidative injury to the gut.<sup>10-11</sup> Deterioration of antioxidative glutathione metabolism<sup>12</sup> and increased colonic oxidative damage to proteins and DNA in association with impaired enzyme activity of CuZn superoxide dismutase has been reported previously in patients with Crohn's disease (CD).<sup>13</sup> Efficacy of current standard therapy in IBD was suggested to be related to antioxidant actions: 5-aminosalicylic acid is a highly potent scavenger of ROS<sup>14</sup> and was shown to reduce lipid peroxidation in CD and ulcerative colitis (UC).<sup>15</sup> Preliminary uncontrolled clinical trials showed that CuZn superoxide dismutase may be of benefit in CD<sup>16</sup> and UC.<sup>17</sup> Several classical antioxidants were reportedly effective in reducing ROS production in UC<sup>18</sup> and experimental colitis.<sup>7-8</sup> Intestinal inflammation induced by administration of an ROS-generating agent in rats could be reversed by supplementation of glutathione.<sup>19</sup> It has therefore been proposed that IBD results from an imbalance between pro-oxidant and antioxidant mechanisms.<sup>20-21</sup>

The tripeptide glutathione (L- $\gamma$ -glutamyl-L-cysteinylglycine, GSH) is the most important intracellular antioxidant. In the presence of ROS reduced GSH (rGSH) is oxidised to glutathione disulphide (GSSG). This reaction is catalysed by GSH peroxidase and reversed by GSH reductase (fig 1).<sup>22</sup> The intestinal GSH content very much depends on GSH synthesis as in vivo administration of a specific inhibitor of  $\gamma$ -glutamylcysteine synthetase ( $\gamma$ GCS) results in a rapid decrease of GSH within 30 minutes, ultimately leading to mucosal damage after several days.<sup>23</sup>  $\gamma$ GCS, the rate limiting enzyme in GSH synthesis (labelled 2 in fig1), is controlled via a feedback mechanism by the end product GSH.<sup>24</sup> High intracellular glutamate levels support GSH synthesis as glutamate competes with GSH at the regulatory site of the enzyme  $\gamma$ GCS.<sup>24</sup> However, high extracellular glutamate concentrations impair GSH

Department of  
Surgery, University of  
Heidelberg,  
Heidelberg, Germany  
B Sido  
A Hochlehnert  
C Herfarth

Department of  
Immunochemistry,  
German Cancer  
Research Center,  
Heidelberg, Germany  
V Hack  
H Lipps  
W Dröge

Correspondence to:  
Dr B Sido, Department of  
Surgery, University of  
Heidelberg, Im Neuenheimer  
Feld 110, 69120 Heidelberg,  
Germany.

Accepted for publication  
31 October 1997






Figure 2 Fasting plasma cysteine levels in patients with Crohn's disease (n=33) and ulcerative colitis (n=33) before surgery and 10 days and three months after complete resection of inflamed bowel. Bars represent mean (SE). Data were statistically analysed by the Student's *t* test for unpaired samples including a Bonferroni's correction. \**p*<0.008, †*p*<10<sup>-4</sup> versus controls (n=65).




Figure 3 Linear regression analysis of preoperative plasma cysteine levels plotted against the individual clinical activity index according to Rachmilewitz in patients with ulcerative colitis.

was confirmed in each series. Pulverised tissue was homogenised in 2.5% SSA by sonification (Branson Sonifier 250, Branson Ultrasonics, Danbury, USA) and centrifuged at 4°C (12 000 *g*, 10 minutes). For analysis of total GSH (tGSH) in mucosa, muscle, PBMC, and RBC the supernatant was assayed enzymatically as previously described with minor modifications.<sup>34</sup> Briefly, 10 µl supernatant of




Figure 4 Total thiol levels in non-deproteinised plasma in patients with Crohn's disease (n=33) and ulcerative colitis (n=33) before surgery and 10 days and three months after complete resection of inflamed bowel. Values represent means (SE) and were statistically analysed by the Student's *t* test for unpaired samples. According to Bonferroni's correction the level of significance was set at *p*<0.008. \**p*<10<sup>-4</sup> versus controls (n=65).

the relevant sample was mixed with 90 µl SSA (2.5%), 6 µl triethanolamine, 0.5 ml buffer (0.1 M Na<sub>2</sub>HPO<sub>4</sub>; 5 mM EDTA; pH 7.5) and 0.2 ml DTNB (3 mM) and 1 U yeast glutathione reductase (Sigma, Deisenhofen, Germany). Finally, 0.2 ml NADPH (1 mM) was added and the absorbance recorded at 41 nm. The rate of reaction at 25°C was expressed as the change in absorbance per six minutes. GSSG was determined according to the method of Griffith<sup>35</sup> after derivatisation of reduced GSH (rGSH) in 0.1 ml supernatant with 2 µl 2-vinylpyridine for 30 minutes at room temperature. Analysis then proceeded as described above. Nine freshly prepared solutions with different concentrations of GSH and GSSG, respectively, were used as standards to calculate the GSH/GSSG concentration in the tissue samples from a linear correlation between absorbance and concentration. The amount of rGSH was calculated as follows: rGSH=tGSH-2 × GSSG. Protein was determined by the Lowry method.<sup>36</sup>

For measurement of enzyme activity of γGCS mucosa samples were sonificated in 0.1 M Tris-HCl buffer (pH 8.0) containing 0.15 M KCl, 20 mM MgCl<sub>2</sub>, and 2 mM Na<sub>2</sub>EDTA. Analysis was performed using a coupled enzyme procedure according to the method of Seelig.<sup>37</sup> Enzyme activity in the reaction mixture was obtained from the decrease in absorbance of NADH at 340 nm (37°C). For measurement of γ-glutamyl transferase (γGT) activity mucosa samples were sonificated in 0.4 ml Tris-HCl buffer (0.1 M, pH 8.25) and analysis was performed using a commercially available assay kit (Boehringer Mannheim, Mannheim, Germany). For both enzymes a linear increase in enzyme activity with increasing volumes of tissue homogenate and a linear correlation between absorbance and reaction time has been confirmed in a separate set of experiments. The protein content in the homogenate was determined according to Bradford's method.<sup>38</sup>

ANALYSIS OF DATA

Group comparison for statistical analysis was performed using the Student's *t* test for unpaired samples (two tailed) after analysis of variance. The level of significance (*p*<0.05) was corrected according to Bonferroni to allow for multiple comparisons. Results are presented as mean (SE).

Results

CONSTITUENT AMINO ACIDS OF GSH IN PLASMA AND INTESTINE

The constituent amino acids cysteine and its disulphide cystine, as well as glutamate and glycine were measured in plasma of patients with CD and UC preoperatively, and 10 days and three months postoperatively after resection of inflamed bowel. In both groups of IBD cysteine was found to be significantly decreased before the operation in the presence of intestinal inflammation compared with healthy control subjects (fig 2). Lowest cysteine levels were observed in patients with UC. Cysteine levels rapidly normalised early after removal of

Table 1 Perioperative plasma concentrations ( $\mu\text{M}$ ) of GSH precursor amino acids in patients with IBD

	Cystine	Glutamate	Glycine
Control (n=65)	61.3 (1.7)	28.0 (1.6)	247 (5)
Crohn's disease (n=33)			
Before surgery	42.8 (2.4)***	32.3 (3.3)	254 (14)
10 days after surgery	56.0 (3.0)	36.2 (2.8)**	324 (20)***
3 months after surgery	52.7 (2.8)**	31.6 (3.0)	268 (15)
Ulcerative colitis (n=33)			
Before surgery	47.3 (1.8)***	36.7 (3.2)**	234 (13)
10 days after surgery	64.3 (2.4)	44.0 (3.1)***	327 (19)***
3 months after surgery	64.5 (3.6)	34.3 (2.5)	337 (23)***

Values are expressed as mean (SE).

\*\* $p < 0.01$ , \*\*\* $p < 0.001$  v healthy controls (n=65) (Student's *t* test for unpaired samples, Bonferroni's correction).

Table 2 Concentration (nmol/mg protein) of GSH precursor amino acids in ileal mucosa from patients with Crohn's disease

	Cystine	Glutamate	Glycine
Controls (n=21)	0.92 (0.13)	39.7 (3.4)	33.0 (2.9)
Crohn's disease			
Non-inflamed (n=21)	0.80 (0.14)	37.1 (2.6)	30.2 (2.0)
Inflamed (n=26)	0.83 (0.16)	33.0 (3.2)	26.1 (2.2)

Values are expressed as mean (SE). Data were analysed using the Student's *t* test for unpaired samples. Significance was set at  $p < 0.025$ , according to Bonferroni's correction.

inflamed bowel and did not differ from controls three months after the operation in both IBD groups. Interestingly, a separate group of patients with CD without intestinal inflammation, but with manifestation of chronic perianal fistulas, had normal cysteine levels (13.9 (0.7)  $\mu\text{M}$  versus 15.3 (0.4)  $\mu\text{M}$  in controls; NS). In patients with UC low preoperative cysteine levels were closely correlated with a high clinical activity index (CAI) of Rachmilewitz ( $r = 0.64$ ,  $p < 10^{-4}$ ; fig 3). Median cysteine levels in controls (CAI 0) and UC patients with low activity (CAI 1–6), moderate activity (CAI 7–12), and high activity (CAI >12) were 15.3, 12.2, 8.9, and 7.9  $\mu\text{M}$ , respectively. Such a correlation could not be established in CD patients using the CDAI index according to Best. Similarly, the total thiol content in deproteinised plasma in IBD, mainly representing protein sulfhydryls, was significantly reduced to the same extent in both CD and UC preoperatively (fig 4). Compared with cysteine, reduction of total thiol levels in plasma persisted early postoperatively but returned to normal values three months after removal of inflamed bowel.

Cystine levels were significantly decreased in both CD and UC patients in the presence of intestinal inflammation (table 1) but rapidly normalised early after the operation. While UC patients maintained high cysteine levels three months after restorative proctocolectomy, plasma cystine concentrations dropped again

below controls in patients with CD. Glycine levels did not differ from controls in both IBD groups before the operation. However, a long lasting increase was recorded in patients with UC after resection of inflamed bowel, whereas glycine levels initially increased early postoperatively in patients with CD and again reached preoperative values three months later. Glutamate, which is known to inhibit intracellular uptake of cystine competitively,<sup>25</sup> was the only constituent amino acid found to be above control levels before the operation. We observed a 22% higher mean concentration of glutamate in patients with CD compared with controls; the 36% increase seen in patients with UC did not completely return to normal at the end of the observation period.

In contrast to the presented variations in plasma levels of precursor amino acids of GSH, mucosal concentrations of cystine, glutamate, and glycine consistently tended to be lower in the ileum of patients with CD by 10%, 17%, and 21%, respectively, compared with controls (table 2). Mucosal cystine concentrations did not parallel the pronounced decrease in plasma levels and mucosal uptake of glutamate was somewhat impaired despite high plasma levels in patients with CD.

#### INTRACELLULAR GSH CONTENT IN PBMC, RBC, AND MUSCLE

Preoperative plasma analysis revealed a pattern of amino acids which suggests an impairment of GSH synthesis. However, analysing PBMC in the same CD patients we were able to demonstrate that intracellular rGSH and GSSG levels did not differ from controls irrespective of inflammation (table 3). Similar values, although slightly lower, were obtained in patients with UC. rGSH levels were even found to be increased in RBC in patients with CD compared with controls (13.8 (0.9) versus 9.6 (0.5) nmol/mg protein;  $p < 10^{-4}$ ). This finding was not associated with intestinal inflammation, as results were identical three months after removal of inflamed bowel (15.0 (1.0) nmol/mg protein;  $p = 10^{-6}$  versus controls). A normal redox status of glutathione in both PBMC and RBC, as defined by the ratio rGSH/GSSG, argues against considerable oxidative stress in the blood compartment in IBD. Similarly, tissue levels of rGSH in the muscle of patients with CD were not decreased despite plasma cyst(e)ine deficiency (11.6 (1.4) versus 13.3 (1.6) nmol/mg protein in controls; NS). As GSSG in muscle was marginally above control values in patients with CD (0.91 (0.16)

Table 3 Glutathione status in peripheral blood mononuclear cells (PBMC) in IBD

	rGSH (nmol/mg protein)	GSSG (nmol/mg protein)	rGSH/GSSG
Control (n=65)	22.8 (1.2)	2.05 (0.11)	12.2 (0.8)
Crohn's disease (n=33)			
Before surgery	23.0 (2.0)	2.06 (0.17)	14.7 (2.0)
10 days after surgery	20.2 (1.4)	2.23 (0.21)	11.8 (1.5)
3 months after surgery	22.2 (1.5)	2.19 (0.20)	12.0 (1.2)
Ulcerative colitis (n=33)			
Before surgery	19.5 (1.3)	2.25 (0.24)	11.5 (1.3)
10 days after surgery	17.9 (1.3)	2.43 (0.25)	10.0 (1.2)
3 months after surgery	20.7 (1.1)	2.30 (0.20)	11.0 (1.1)

Values are expressed as mean (SE). Data were analysed using the Student's *t* test for unpaired samples. Significance was set at  $p < 0.008$ , according to Bonferroni's correction.




Figure 5 Pro-oxidant glutathione status in the mucosa of freshly resected ileum in patients with Crohn's disease. Reduced glutathione (rGSH) is decreased in inflamed ( $n=26$ ) as well as non-inflamed ( $n=21$ ) mucosa in Crohn's disease. Oxidised glutathione (GSSG) is increased in areas of inflammation so that the redox status of glutathione, as defined by the ratio rGSH/GSSG, is heavily decreased in inflamed mucosa. Bars represent mean (SE). Statistical analysis was performed by the Student's *t* test including a Bonferroni's correction. \*\*\* $p<0.01$ , † $p<10^{-4}$  versus controls ( $n=21$ ).

versus 0.74 (0.16) nmol/mg protein; NS), the glutathione redox status tended to be decreased in this tissue but did not reach statistical significance (17.1 (2.3) versus 23.7 (2.9) in controls). In summary, deficiency of the sulphur-containing precursor amino acids of GSH in plasma does not result in an impairment of GSH synthesis in blood cells and muscle in IBD, irrespective of the presence or absence of intestinal inflammation.

#### GUT MUCOSAL GSH CONTENT

The finding of unimpaired GSH synthesis in the blood cells and muscle prompted us to investigate the GSH metabolism in inflamed and non-inflamed mucosa of the small bowel. Only patients with CD were considered for

these investigations because the large bowel was subject to long ischaemia times during proctocolectomy in patients with UC. In contrast to results obtained in blood cells and muscle, mucosal rGSH was profoundly reduced by 24% and 38% in non-inflamed and inflamed mucosa, respectively (fig 5). Despite the shortage of rGSH, GSSG significantly increased with inflammation. Consequently, the GSH redox status (ratio rGSH/GSSG) dramatically decreased down to 42% in the inflamed ileum. This difference between patients with CD and controls is not simply the consequence of chronic steroid therapy as the GSH redox status in inflamed mucosa of patients receiving steroids ( $n=13$ ) was 18.1 (1.8) versus 16.0 (1.7) in patients without steroids ( $n=13$ ) (NS).




Figure 6 Decreased mucosal enzyme activity of gamma-glutamylcysteine synthetase ( $\gamma$ GCS) in inflamed ( $n=26$ ) and non-inflamed ( $n=21$ ) ileum of patients with Crohn's disease. Bars represent mean (SE). Statistical analysis was performed by the Student's *t* test for unpaired samples including a Bonferroni's correction. \* $p<0.025$ , † $p<10^{-4}$  versus controls ( $n=21$ ).

#### GUT MUCOSAL $\gamma$ GCS ACTIVITY

$\gamma$ GCS is the rate limiting enzyme in GSH synthesis and activity is negatively regulated via a feedback mechanism by the amount of GSH produced.<sup>24</sup> In CD, however, we observed a profound decrease in  $\gamma$ GCS enzyme activity despite GSH deficiency in inflamed and non-inflamed small bowel mucosa (fig 6). Chronic steroid therapy did not improve  $\gamma$ GCS enzyme activity which was 95.2 (12.1) nmol/min/mg protein in patients receiving steroids and 83.2 (5.6) nmol/min/mg protein in patients not receiving steroids (NS).

#### GUT MUCOSAL $\gamma$ GT ACTIVITY

$\gamma$ GT is located at the luminal side of the brush border membrane of intestinal epithelial cells. It degrades luminal GSH and GSSG to  $\gamma$ -glutamyl amino acids that are transported to the cell where they are ultimately used for resynthesis of GSH (fig 1).<sup>28</sup> Similarly to  $\gamma$ GCS,  $\gamma$ GT activity was considerably decreased in inflamed mucosa of the ileum compared with controls and was 17.7 (1.4) and 16.7 (1.6) mU/mg protein in patients with and




Figure 7 Decreased mucosal enzyme activity of  $\gamma$ -glutamyl transferase ( $\gamma$ GT) in inflamed ( $n=26$ ) and non-inflamed ( $n=21$ ) ileum of patients with Crohn's disease. Bars represent mean (SE). Statistical analysis was performed by the Student's  $t$  test for unpaired samples. According to Bonferroni's correction the level of significance was set at  $p < 0.025$ . \* $p < 10^{-4}$  versus controls ( $n=21$ ).

without steroids, respectively (NS). In non-inflamed mucosa of patients with CD enzyme activity was intermediate between controls and inflamed tissue (fig 7).

### Discussion

In this study we present a detailed analysis of gut mucosal GSH synthesis in patients with CD. Decreased activities of enzymes involved in GSH synthesis ( $\gamma$ GCS,  $\gamma$ GT) predominantly contribute to intestinal GSH deficiency in both non-inflamed and inflamed ileum. Because the GSH level was shown to be normal or even increased in extraintestinal compartments, decreased availability of the sulphur-containing constituent amino acids of GSH in plasma (but not in the mucosa) seems to be of minor importance. This impaired mucosal antioxidative capacity ultimately increases gut susceptibility to oxidative tissue injury. Despite intestinal rGSH deficiency, the GSSG level increased with inflammation resulting in a pro-oxidant shift of the mucosal redox balance of GSH (rGSH/GSSG) in CD. This could have major implications on intestinal immune functions in IBD as pro-oxidant mechanisms are known to be involved in the activation of the transcription factor NF $\kappa$ B, thereby mediating the transcription of several immunologically relevant cytokines and cell adhesion molecules.<sup>39</sup>

Cysteine and its disulphide cystine are known to be limiting substrates in GSH biosynthesis<sup>22,27</sup> and therefore generate a strong influence on intracellular GSH stores.<sup>33,34,40</sup> Reduced plasma cysteine and cystine levels are not unique to patients with inflammatory bowel disease and could also be demonstrated in other catabolic diseases such as HIV infection<sup>33</sup> and advanced malignant tumour growth<sup>41,42</sup> in conjunction with a reduction in intracellular GSH levels in PBMC. In patients

with IBD cysteine deficiency seems to be a consequence and not a primary mechanism of relapsing inflammation because (1) cysteine and cystine levels rapidly normalise after removal of inflamed bowel in both CD and UC, (2) patients with CD without intestinal inflammation but with manifestations of chronic perianal fistulas have normal cysteine values, (3) UC, which usually involves larger bowel segments, is accompanied by lower cysteine levels than CD, and (4) preoperative cysteine levels significantly correlate with the clinical disease activity in patients with UC. Plasma cysteine levels could be decreased due to ongoing inflammation for several reasons. The sulphhydryl amino acid cysteine is a potent scavenger of hydroxyl radicals and could be readily consumed due to oxidative stress in the inflamed gut. In support of this explanation is our finding that the decrease in total thiol content in plasma, mainly representing protein sulphhydryls, is also associated with inflammation in both CD and UC. Alternatively, hypochlorite that is produced from hydrogen peroxide and chloride via the myeloperoxidase reaction in neutrophils was shown to be present in the mucosa of rectal biopsy samples in patients with IBD.<sup>43</sup> The reaction of hypochlorite with ammonia yields monochloramine. Both hypochlorite and monochloramine avidly degrade cysteine<sup>44</sup> and could account for cysteine deficiency. Inversely, the therapeutically effective drug 5-aminosalicylic acid is a potent antioxidant,<sup>44</sup> inhibits neutrophil production of hypochlorite and monochloramine,<sup>44-46</sup> and protects cysteine from degradation by these two toxic compounds.<sup>44</sup> Supplementation of cysteine or N-acetyl-L-cysteine might have beneficial effects on disease activity due to its radical scavenging activity and as it helps to replenish intracellular GSH stores. Oral application of cysteine has recently been shown to ameliorate tissue injury in experimental colitis.<sup>47</sup>

Although the pattern of precursor amino acid, as described in the present study, suggests impairment of GSH synthesis in IBD, our results do not argue for overall inhibition of GSH synthesis. Among the compartments analysed we could show a decrease of GSH in the intestine, but not in blood cells and muscle. The reason for the increased GSH content in RBC is presently not known but has also been observed in tumour bearing mice in the presence of decreased plasma cyst(e)ine levels.<sup>48</sup> It is unlikely that plasma deficiency of cyst(e)ine is due to avid amino acid uptake into RBC for subsequent GSH synthesis as plasma cyst(e)ine levels normalise postoperatively, although increased GSH levels persist in RBC. In cultured cells cystine is taken up by the membrane amino acid transport system  $x_c^-$  that imports cystine in exchange for glutamate.<sup>49</sup> In vitro this transport system can be induced by glutathione deficiency or oxidative stress.<sup>50</sup> Provided that such a mechanism is active in vivo, induction of the transport system could explain why mucosal levels of cystine were not found to be decreased to the same extent below control levels as were plasma levels and,

inversely, why glutamate levels were decreased in the ileum. In the face of a normal or even increased GSH concentration in extraintestinal compartments, decreased mucosal glutathione levels are not solely due to insufficient uptake of precursor amino acids from plasma. Rather, it has to be assumed that intestinal GSH synthesis is impaired. The gut is highly dependent on GSH synthesis as has been shown in mice. Treatment with L-buthionine-SR-sulphoximine, a specific inhibitor of  $\gamma$ GCS, rapidly decreased mucosal content of GSH down to 60% of control values within 30 minutes.<sup>23</sup> In the present study we could demonstrate a 45% downregulation of mucosal  $\gamma$ GCS activity in patients with CD that could easily account for decreased GSH levels in the gut mucosa.  $\gamma$ GCS, the rate limiting enzyme in GSH synthesis, is negatively regulated via a feedback mechanism by which high concentrations of the end product GSH inhibit the enzyme.<sup>24</sup> Obviously, the physiological feedback regulation of  $\gamma$ GCS by GSH is not operative in patients with CD despite mucosal GSH deficiency. Feedback inhibition of  $\gamma$ GCS by GSH can be blocked by a high concentration of intracellular glutamate that competes with GSH at the regulatory site of the enzyme.<sup>24</sup> However, such a mechanism can also be excluded as mucosal glutamate levels were shown to be below controls in CD.

Loss of intracellular GSH could also cause mucosal GSH deficiency. GSSG formed by oxidative stress is either reduced to GSH via the glutathione reductase-catalysed reaction or, when produced in excess amounts, is transported from the cells to the blood in order to maintain equilibrium levels of GSH and GSSG.<sup>50</sup> Alternatively, GSSG reacts with protein sulfhydryls via a mixed disulphide reaction.<sup>51</sup> The net result of both mechanisms is a loss of intracellular free GSH. Another drain on the intracellular pool of GSH is the glutathione S-transferase reaction which catalyses the conjugation of glutathione with a large number of electrophilic compounds. Although glutathione S-transferase was reported to be present in the intestine,<sup>52</sup> our laboratory could provide evidence for a decrease in glutathione S-transferase activity in the gut mucosa in CD (unpublished observation).

Intestinal epithelial cells were shown to contain high levels of  $\gamma$ GT activity predominantly at the villous tips.<sup>53</sup> We could demonstrate a significant decrease in  $\gamma$ GT activity in the inflamed ileum of patients with CD. Consequently, degradation products of luminal GSH cannot be made available to the intestine for resynthesis of GSH to maintain physiological GSH levels. However, previous studies in vascularly perfused rat small intestine and in vivo studies in rats with specific inhibition of  $\gamma$ GT by acivicin showed that the salvage pathway via  $\gamma$ GT is of minor importance.<sup>12, 54</sup> This finding is supported by studies from our laboratory which have shown that the decrease in colonic GSH levels in CD is identical to the ileum, although  $\gamma$ GT activity in the colon was found to be only 31% of the activity in the ileum and remains unchanged in CD (unpub-

lished data). Thus the present study provides evidence that several factors relevant to GSH synthesis in gut mucosa are impaired; the reduced blood supply of the precursor amino acids cysteine and cystine in conjunction with downregulation of mucosal  $\gamma$ GT activity and, most important, profound reduction in  $\gamma$ GCS enzyme activity argue for insufficient GSH synthesis as a primary mechanism for mucosal GSH deficiency.

GSH is essential for the function and structural integrity of the gut; GSH-deficient mice show severe degeneration of the jejunum and colonic mucosa and were found to have body weight loss and diarrhoea.<sup>23</sup> Oral administration of a mixture of constituent amino acids (L-glutamate, L-cysteine, L-glycine) was shown not to be effective in increasing gut mucosal GSH levels.<sup>12, 23</sup> In contrast, high amounts of GSH result in supply of GSH to the lower gastrointestinal tract and, probably due to uptake of intact GSH from the lumen across the brush border membrane, thus provide a direct means of raising GSH levels in the mucosa of the jejunum, ileum and colon.<sup>12, 23, 30</sup> Based on the profound impairment of GSH synthesis as demonstrated in the present study, GSH supplementation might be of therapeutic value in IBD.

The study was financially supported by the DKFZ project 072006.

- Mahida YR, Wu KC, Jewell DP. Respiratory burst activity of intestinal macrophages in normal and inflammatory bowel disease. *Gut* 1989;30:1362-70.
- Williams JG. Phagocytes, toxic oxygen metabolites and inflammatory bowel disease: implications for treatment. *Ann R Coll Surg Engl* 1990;72:253-62.
- Grisham MB, Granger DN. Neutrophil-mediated mucosal injury. Role of reactive oxygen metabolites. *Dig Dis Sci* 1988;33:65-15S.
- Weiss SJ. Tissue destruction by neutrophils. *N Engl J Med* 1989;320:365-76.
- Williams JG, Hughes LE, Hallert MB. Toxic oxygen metabolite production by circulating phagocytic cells in inflammatory bowel disease. *Gut* 1990;31:187-93.
- Grisham MB, Volkmer C, Tso P, et al. Metabolism of trinitrobenzene sulfonic acid by rat colon produces reactive oxygen species. *Gastroenterology* 1991;101:540-7.
- Millar AD, Rampton DS, Chander CL, et al. Evaluating the antioxidant potential of new treatments for inflammatory bowel disease using a rat model of colitis. *Gut* 1996;39:407-15.
- Keshavarazian A, Sedghi S, Kanovsky J, et al. Excessive production of reactive oxygen metabolites by inflamed colon: analysis by chemiluminescence probe. *Gastroenterology* 1992;103:177-85.
- Grisham MB, MacDermott RP, Deitch EA. Oxidant defense mechanisms in the human colon. *Inflammation* 1990;6:669-80.
- Babbs C. Oxygen radicals in ulcerative colitis. *Free Radic Biol Med* 1992;13:169-81.
- van der Vliet A, Bast A. Role of reactive oxygen species in intestinal diseases. *Free Radic Biol Med* 1992;12:499-513.
- Iantomasi T, Marraccini P, Favilli F, et al. Glutathione metabolism in Crohn's disease. *Biochem Med Metabol Biol* 1994;53:87-91.
- Lih-Brody L, Powell SR, Collier KP, et al. Increased oxidative stress and decreased antioxidant defenses in mucosa of inflammatory bowel disease. *Dig Dis Sci* 1996;41:2078-86.
- Ahnfelt-Ronne I, Nielsen OH. The anti-inflammatory moiety of sulfasalazine, 5-aminosalicylic acid, is a radical scavenger. *Agents Actions* 1987;21:191-4.
- Ahnfelt-Ronne I, Nielsen OH, Christensen A, et al. Clinical evidence supporting the radical scavenger mechanism of 5-aminosalicylic acid. *Gastroenterology* 1990;98:1162-9.
- Emerit J, Pelletier S, Likforman J, et al. Phase II trial of superoxide dismutase (CuZn SOD) in the treatment of Crohn's disease. *Free Radic Res Commun* 1991;12-13(Pt 2):563-9.
- Niwa Y, Somiyai K, Michelson AM, et al. Effect of liposomal-encapsulated superoxide dismutase on active oxygen-related disorders. *Free Radic Res Commun* 1985;1:137-53.
- Simmonds NJ, Blake DR, Rampton DS. The effects of aminosalicylates and potential new drugs on mucosal reactive oxygen metabolite production in ulcerative colitis. *Gut* 1992;33(suppl):S7.

- 19 Tamai H, Levin S, Gagarella TS. Induction of colitis in rats by 2-2'-azobis(2-amidinopropane) dihydrochloride. *Inflammation* 1992;16:69-81.
- 20 Grisham MB. Oxidants and free radicals in inflammatory bowel disease. *Lancet* 1994;344:859-61.
- 21 Gross V, Arndt H, Andus T, et al. Free radicals in inflammatory bowel diseases. Pathophysiology and therapeutic implications. *HepatoGastroenterol* 1994;41:320-7.
- 22 Meister A, Anderson ME. Glutathione. *Ann Rev Biochem* 1983;52:711-60.
- 23 Martensson J, Jain A, Meister A. Glutathione is required for intestinal function. *Proc Natl Acad Sci USA* 1990;87:1715-9.
- 24 Richman P, Meister A. Regulation of gamma-glutamyl-cysteine synthetase by nonallosteric feedback inhibition by glutathione. *Biol Chem* 1975;250:1422-6.
- 25 Eck H-P, Dröge W. Influence of the extracellular glutamate concentration on the intracellular cyst(e)ine concentration in macrophages and on the capacity to release cysteine. *Biol Chem Hoppe-Seyler* 1989;370:109-13.
- 26 Bannai S, Kitamura E. Transport interaction of L-cystine and L-glutamate in human diploid fibroblasts in culture. *J Biol Chem* 1980;255:2372-6.
- 27 Meister A. Selective modification of glutathione metabolism. *Science* 1983;220:472-7.
- 28 Vincenzini MT, Favilli F, Iantomasi T. Intestinal uptake and transmembrane transport systems of intact GSH: characteristics and possible biological role. *Biochim Biophys Acta* 1992;1113:13-23.
- 29 Hagen TM, Wierzbicka GT, Sillau AH, et al. Bioavailability of dietary glutathione: effect on plasma concentrations. *Am J Physiol* 1990;259:G524-9.
- 30 Hagen TM, Wierzbicka GT, Bowman BB, et al. Fate of dietary glutathione: disposition in the gastrointestinal tract. *Am J Physiol* 1990;259:G530-5.
- 31 Best WR, Becktel JM, Singleton JW, et al. Development of a Crohn's disease activity index. National Cooperative Crohn's Disease Study. *Gastroenterology* 1976;70:439-44.
- 32 Rachmilewitz D. Coated mesalazine (5-aminosalicylic acid) versus sulphasalazine in the treatment of active ulcerative colitis: a randomised trial. *BMJ* 1989;298:82-6.
- 33 Eck H-P, Gmünder H, Hartmann M, et al. Low concentrations of acid-soluble thiol (cysteine) in the blood plasma of HIV-1-infected patients. *Biol Chem Hoppe-Seyler* 1989;370:101-8.
- 34 Gmünder H, Eck HP, Benninghoff B, et al. Macrophages regulate intracellular glutathione levels of lymphocytes. Evidence for an immunoregulatory role of cysteine. *Cell Immunol* 1990;129:32-46.
- 35 Griffith OW. Determination of glutathione and glutathione disulfide using glutathione reductase and 2-vinyl-pyridine. *Anal Biochem* 1980;106:207-12.
- 36 Lowry OH, Rosenbrough WJ, Farr AL, et al. Protein measurement with the folin phenol reagent. *J Biol Chem* 1951;193:265-75.
- 37 Seelig GF, Meister A. Glutathione biosynthesis;  $\gamma$ -glutamylcysteine synthetase from rat kidney. *Methods Enzymol* 1985;13:379-90.
- 38 Bradford MM. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248-54.
- 39 Siebenlist U, Franzoso G, Brown K. Structure, regulation and function of NF- $\kappa$ B. *Annu Rev Cell Biol* 1994;10:405-55.
- 40 Ishii T, Sugita Y, Bannai S. Regulation of glutathione in mouse spleen lymphocytes by transport of cysteine. *J Cell Physiol* 1987;133:330-6.
- 41 Eck H-P, Drings P, Dröge W. Plasma glutamate levels, lymphocyte reactivity and death rate in patients with bronchial carcinoma. *J Cancer Res Clin Oncol* 1989;115:571-4.
- 42 Hack V, Schmid D, Breikreutz R, et al. Cystine levels, cystine flux and protein catabolism in cancer cachexia, HIV/SIV infection and senescence. *FASEB J* 1997;11:84-92.
- 43 Simmonds NJ, Allen RE, Stevens TR, et al. Chemiluminescence assay of mucosal reactive oxygen metabolites in inflammatory bowel disease. *Gastroenterology* 1992;103:186-96.
- 44 Tamai H, Kachur JF, Grisham MB, et al. Scavenging effect of 5-aminosalicylic acid on neutrophil-derived oxidants. Possible contribution to the mechanism of action in inflammatory bowel disease. *Biochem Pharmacol* 1991;41:1001-6.
- 45 Williams JG, Hallett MB. The reaction of 5-amino-salicylic acid with hypochlorite. Implications for the mode of action in inflammatory bowel disease. *Biochem Pharmacol* 1989;38:149-54.
- 46 Dallegrì F, Ottonello L, Ballestrero A, et al. Cytoprotection against neutrophil derived hypochlorous acid: a potential mechanism for the therapeutic action of 5-aminosalicylic acid in ulcerative colitis. *Gut* 1990;31:184-6.
- 47 Karmeli F, Rachmilewitz D. Cysteine and methionine: a novel modality to ameliorate tissue injury in experimental colitis [abstract]. *Gastroenterology* 1997;112:A1009.
- 48 Hack V, Gross A, Kinscherf R, et al. Abnormal glutathione and sulfate levels after interleukin 6 treatment and in tumor-induced cachexia. *FASEB J* 1996;10:1219-26.
- 49 Bannai S. Exchange of cystine and glutamate across plasma membrane of human fibroblasts. *J Biol Chem* 1986;261:2256-62.
- 50 Deneke SM, Fanburg BL. Regulation of cellular glutathione. *Am J Physiol* 1989;257:L163-73.
- 51 DeLucia AJ, Mustafa MG, Hussain MZ, et al. Ozone interaction with rodent lung. III. Oxidation of reduced glutathione and formation of mixed disulfides between protein and nonprotein sulfhydryls. *J Clin Invest* 1975;55:794-802.
- 52 Peters WHM, Roelofs MMJ, Nagengast FM, et al. Human intestinal glutathione S-transferases. *Biochem J* 1989;257:471-6.
- 53 Cornell JS, Meister A. Glutathione and  $\gamma$ -glutamyl cycle enzymes in crypt and villus tip cells of rat jejunal mucosa. *Proc Natl Acad Sci USA* 1976;73:420-2.
- 54 Hagen TM, Jones DP. Transepithelial transport of glutathione in vasculature perfused small intestine of rat. *Am J Physiol* 1987;252:G607-13.